


ENSO: Recent Evolution, Current Status and Predictions



Update prepared by:
Climate Prediction Center / NCEP
30 October 2017

Outline

Summary

Recent Evolution and Current Conditions

Oceanic Niño Index (ONI)

Pacific SST Outlook

U.S. Seasonal Precipitation and Temperature Outlooks

Summary

Summary

ENSO Alert System Status: La Niña Watch

ENSO-Neutral conditions are present.*

Equatorial sea surface temperatures (SSTs) are near-to-below average across the central and eastern Pacific Ocean.

La Niña conditions are favored (~55%-65%) during the Northern Hemisphere fall and winter 2017-18.*

* Note: These statements are updated once a month (2nd Thursday of each month) in association with the ENSO Diagnostics Discussion, which can be found by clicking [here](#).

Recent Evolution of Equatorial Pacific SST Departures (°C)


Below average SSTs spanned most of the central and eastern Pacific Ocean through January 2017.

During January and February 2017, above-average SSTs expanded in the eastern Pacific Ocean.

From mid April to July 2017, near-to-above average SSTs spanned most of the equatorial Pacific.

During August 2017, above-average SSTs dissipated east of the date line.



Since September, SSTs have been near-to-below average across the central and eastern Pacific.



Niño Region SST Departures (°C) Recent Evolution


The latest weekly SST departures are:

Niño 4	-0.2°C
Niño 3.4	-0.5°C
Niño 3	-0.8°C
Niño 1+2	-1.4°C



SST Departures (°C) in the Tropical Pacific During the Last Four Weeks


During the last four weeks, equatorial SSTs were below average across the east-central and eastern Pacific Ocean, and above average in the far western Pacific.



Weekly SST Departures during the Last Four Weeks


During the last four weeks, below-average SSTs persisted across the eastern Pacific Ocean. SST anomalies were more variable in the central and east-central Pacific.

Weekly SST Anomalies (DEG C)



Change in Weekly SST Departures over the Last Four Weeks

During the last four weeks, changes in equatorial SST anomalies were mostly negative across the eastern Pacific Ocean.






Upper-Ocean Conditions in the Equatorial Pacific

The basin-wide equatorial upper ocean (0-300 m) heat content is greatest prior to and during the early stages of a Pacific warm (El Niño) episode (compare top 2 panels), and least prior to and during the early stages of a cold (La Niña) episode.

The slope of the oceanic thermocline is least (greatest) during warm (cold) episodes.


Recent values of the upper-ocean heat anomalies (below average) and thermocline slope index (near average) reflect ENSO-Neutral conditions.

The monthly thermocline slope index represents the difference in anomalous depth of the 20°C isotherm between the western Pacific (160°E-150°W) and the eastern Pacific (90°-140°W).




Central and Eastern Pacific Upper-Ocean (0-300 m) Weekly Average Temperature Anomalies

Negative subsurface temperature anomalies were present through December 2016. Positive anomalies were present from mid-January through March 2017 before weakening to near zero. Starting in mid-April and mid-June, positive anomalies strengthened before tapering off again. Since mid-July, anomalies have decreased and are negative.




Sub-Surface Temperature Departures in the Equatorial Pacific

In the last two months, negative subsurface temperature anomalies have expanded across the Pacific Ocean.



Recently, the strongest negative anomalies are between 170°W-80°W.




Tropical OLR and Wind Anomalies During the Last 30 Days

Negative OLR anomalies (enhanced convection and precipitation) were evident near the Philippines. Positive OLR anomalies (reduced convection and precipitation) were present over the central Pacific Ocean.

Low-level (850-hPa) winds were near average across most of the tropical Pacific Ocean.

Upper-level (200-hPa) winds were anomalous westerly across most of the tropical Pacific Ocean.



Intraseasonal Variability

Intraseasonal variability in the atmosphere (wind and pressure), which is often related to the Madden-Julian Oscillation (MJO), can significantly impact surface and subsurface conditions across the Pacific Ocean.

Related to this activity:


Significant weakening of the low-level easterly winds usually initiates an eastward-propagating oceanic Kelvin wave.

Weekly Heat Content Evolution in the Equatorial Pacific

From February 2017 through May 2017, positive subsurface temperature anomalies persisted in the western and eastern Pacific Ocean, with oceanic Kelvin waves resulting in anomalous temperature variability in the central Pacific.

During August 2017, an upwelling Kelvin wave resulted in below-average sub-surface temperatures across the east-central and eastern equatorial Pacific. Since September 2017, negative sub-surface anomalies have persisted in those regions.

Equatorial oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Down-welling and warming occur in the leading portion of a Kelvin wave, and up-welling and cooling occur in the trailing portion.



Low-level (850-hPa) Zonal (east-west) Wind Anomalies (m s^{-1})


Low-level easterly wind anomalies generally have persisted over the central and western equatorial Pacific since May 2017.

Since January 2017, westerly wind anomalies were generally observed over the eastern Pacific Ocean.

Since mid-October 2017, the Madden Julian Oscillation (MJO) has contributed to the eastward shift of low-level wind anomalies.

Westerly Wind Anomalies (orange/red shading)

Easterly Wind Anomalies (blue shading)



Upper-level (200-hPa) Velocity Potential Anomalies


Since at least April 2017, anomalous upper-level divergence (green shading) generally persisted near Indonesia, while anomalous convergence (brown shading) persisted near the Date Line.

Eastward propagation of regions of upper-level divergence (green shading) and convergence (brown shading) is particularly evident during January-February 2017, April-May 2017, and since mid-July 2017.

Unfavorable for precipitation (brown shading)

Favorable for precipitation (green shading)

Note: Eastward propagation is not necessarily indicative of the Madden-Julian Oscillation (MJO).




Outgoing Longwave Radiation (OLR) Anomalies

From mid-May to late July 2017, OLR anomalies were negative near Indonesia.

Since mid-August 2017, positive OLR anomalies have persisted over the central Pacific Ocean.

Drier-than-average Conditions (orange/red shading)

Wetter-than-average Conditions (blue shading)



Oceanic Niño Index (ONI)

The ONI is based on SST departures from average in the Niño 3.4 region, and is a principal measure for monitoring, assessing, and predicting ENSO.

Defined as the three-month running-mean SST departures in the Niño 3.4 region. Departures are based on a set of improved homogeneous historical SST analyses (Extended Reconstructed SST - ERSST.v5). The SST reconstruction methodology is described in Huang et al., 2017, J. Climate, vol. 30, 8179-8205.)

It is one index that helps to place current events into a historical perspective

NOAA Operational Definitions for El Niño and La Niña

El Niño: characterized by a positive ONI greater than or equal to $+0.5^{\circ}\text{C}$.

La Niña: characterized by a negative ONI less than or equal to -0.5°C .


By historical standards, to be classified as a full-fledged El Niño or La Niña episode, these thresholds must be exceeded for a period of at least 5 consecutive overlapping 3-month seasons.

CPC considers El Niño or La Niña conditions to occur when the monthly Niño3.4 OISST departures meet or exceed $\pm 0.5^{\circ}\text{C}$ along with consistent atmospheric features. These anomalies must also be forecasted to persist for 3 consecutive months.

ONI (°C): Evolution since 1950

The most recent ONI value (July - September 2017) is -0.1°C .

El Niño ↑
Neutral
La Niña ↓



Historical El Niño and La Niña Episodes Based on the ONI computed using ERSST.v5

Recent Pacific warm (red) and cold (blue) periods based on a threshold of +/- 0.5 °C for the Oceanic Niño Index (ONI) [3 month running mean of ERSST.v5 SST anomalies in the Niño 3.4 region (5N-5S, 120-170W)]. For historical purposes, periods of below and above normal SSTs are colored in blue and red when the threshold is met for a minimum of 5 consecutive over-lapping seasons.


The ONI is one measure of the El Niño-Southern Oscillation, and other indices can confirm whether features consistent with a coupled ocean-atmosphere phenomenon accompanied these periods. The complete table going back to DJF 1950 can be found [here](#).

Year	DJF	JFM	FMA	MAM	AMJ	MJJ	JJA	JAS	ASO	SON	OND	NDJ
2005	0.6	0.6	0.4	0.4	0.3	0.1	-0.1	-0.1	-0.1	-0.3	-0.6	-0.8
2006	-0.8	-0.7	-0.5	-0.3	0.0	0.0	0.1	0.3	0.5	0.7	0.9	0.9
2007	0.7	0.3	0.0	-0.2	-0.3	-0.4	-0.5	-0.8	-1.1	-1.4	-1.5	-1.6
2008	-1.6	-1.4	-1.2	-0.9	-0.8	-0.5	-0.4	-0.3	-0.3	-0.4	-0.6	-0.7
2009	-0.8	-0.7	-0.5	-0.2	0.1	0.4	0.5	0.5	0.7	1.0	1.3	1.6
2010	1.5	1.3	0.9	0.4	-0.1	-0.6	-1.0	-1.4	-1.6	-1.7	-1.7	-1.6
2011	-1.4	-1.1	-0.8	-0.6	-0.5	-0.4	-0.5	-0.7	-0.9	-1.1	-1.1	-1.0
2012	-0.8	-0.6	-0.5	-0.4	-0.2	0.1	0.3	0.3	0.3	0.2	0.0	-0.2
2013	-0.4	-0.3	-0.2	-0.2	-0.3	-0.3	-0.4	-0.4	-0.3	-0.2	-0.2	-0.3
2014	-0.4	-0.4	-0.2	0.1	0.3	0.2	0.1	0.0	0.2	0.4	0.6	0.7
2015	0.6	0.6	0.6	0.8	1.0	1.2	1.5	1.8	2.1	2.4	2.5	2.6
2016	2.5	2.2	1.7	1.0	0.5	0.0	-0.3	-0.6	-0.7	-0.7	-0.7	-0.6
2017	-0.3	-0.1	0.1	0.3	0.4	0.4	0.1	-0.1				

CPC/IRI Probabilistic ENSO Outlook

Updated: 12 October 2017

La Niña is favored (~55%-65%) during the Northern Hemisphere fall and winter 2017-18.



IRI/CPC Pacific Niño 3.4 SST Model Outlook

The multi-model averages predict La Niña to develop and persist through the remainder of the year and into early 2018.





Figure provided by the International Research Institute (IRI) for Climate and Society (updated 17 October 2017).

SST Outlook: NCEP CFS.v2 Forecast (PDF corrected)

Issued: 30 October 2017


The CFS.v2 ensemble mean (black dashed line) favors La Niña through January-March 2018.

CFSv2 forecast Nino3.4 SST anomalies (K) (PDF corrected)



— Latest 8 forecast members
— Earliest 8 forecast members
— Other forecast members
- - - Forecast ensemble mean
— NCDP daily analysis


(Model bias correct base period: 1999–2010; Climatology base period: 1982–2010)



Atmospheric anomalies over the North Pacific and North America During the Last 60 Days

During late August to early September 2017, an anomalous trough (and below-average temperatures) was present over eastern North America, and an anomalous ridge (and above-average temperatures) was present over the western contiguous U.S.


After mid September, the pattern of anomalies flipped. An amplified ridge (and above-average temperatures) occurred over the eastern U.S. On occasion, an anomalous trough (and below-average temperatures) was observed over the western U.S.



Atmospheric anomalies over the North Pacific and North America During the Last 60 Days

During late August to early September 2017, an anomalous trough (and below-average temperatures) was present over eastern North America, and an anomalous ridge (and above-average temperatures) was present over the western contiguous U.S.


After mid September, the pattern of anomalies flipped. An amplified ridge (and above-average temperatures) occurred over the eastern U.S. On occasion, an anomalous trough (and below-average temperatures) was observed over the western U.S.



Atmospheric anomalies over the North Pacific and North America During the Last 60 Days

During late August to early September 2017, an anomalous trough (and below-average temperatures) was present over eastern North America, and an anomalous ridge (and above-average temperatures) was present over the western contiguous U.S.


After mid September, the pattern of anomalies flipped. An amplified ridge (and above-average temperatures) occurred over the eastern U.S. On occasion, an anomalous trough (and below-average temperatures) was observed over the western U.S.




U.S. Temperature and Precipitation Departures During the Last 30 Days

End Date: 28 October 2017

Percent of Average Precipitation




Temperature Departures (degree C)




U.S. Temperature and Precipitation Departures During the Last 90 Days

End Date: 28 October 2017

Percent of Average Precipitation




Temperature Departures (degree C)



U. S. Seasonal Outlooks

November 2017- January 2018

The seasonal outlooks combine the effects of long-term trends, soil moisture, and, when appropriate, ENSO.



Summary

ENSO Alert System Status: La Niña Watch

ENSO-Neutral conditions are present.*

Equatorial sea surface temperatures (SSTs) are near-to-below average across the central and eastern Pacific Ocean.

La Niña conditions are favored (~55%-65%) during the Northern Hemisphere fall and winter 2017-18.*

* Note: These statements are updated once a month (2nd Thursday of each month) in association with the ENSO Diagnostics Discussion, which can be found by clicking [here](#).