

Weather Regime-Dependent Predictability: Sequentially Linked High-Impact Weather Events over the United States during March 2016

Andrew C. Winters*, Lance F. Bosart*, and Daniel Keyser*

*Department of Atmospheric and Environmental Sciences, University at Albany, SUNY

contact: acwinters@albany.edu
This work is funded by NOAA under Grant NA15NWS4680006

Motivation

- High-impact weather events (HWEs), defined by episodes of excessive precipitation or anomalous temperature, can pose important predictability challenges on medium-range (8–16 day) time scales.
- This study introduces the North Pacific Jet (NPJ) phase diagram as a tool to characterize the large-scale flow pattern over the North Pacific.
- The NPJ phase diagram is applied to a period in late March and early April 2016 that was characterized by three sequentially linked HWEs:
 - The 23–24 March 2016 Colorado Front Range snowstorm and Southern Plains severe weather outbreak.
 - The 27–28 March 2016 Ohio River Valley severe weather outbreak.
 - The 3–4 April 2016 Northeast U.S. cold air outbreak.

Data

- 1.0° x 1.0° NCEP Global Forecast System (GFS) and Global Ensemble Forecast System (GEFS) analyses (available every 6 h).
- 0.5° x 0.5° NCEP Climate Forecast System Reanalysis (CFSR; available every 6 h) during 1979–2014.

Conclusions

- The predictability challenges that characterized the 23–24 March 2016 Colorado snowstorm were magnified because uncertainties in the positions of key weather features were concentrated near the Continental Divide.
- Forecasts of the 27–28 March 2016 Ohio River Valley severe weather outbreak exhibited uncertainty due to model disagreement on the placement of a subsynoptic-scale trough over the MS River Valley.
- The evolution of the flow pattern prior to the onset of cold over the Northeast U.S. on 3 April 2016 is consistent with an antecedent environment conducive to extreme cold east of the Rocky Mountains.
- Predictability horizons can vary greatly as a function of weather regime and season. Consequently, regime-dependent predictability, and how it may vary seasonally, is an important science and operations problem worthy of further consideration.

North Pacific Jet Phase Diagram

- The large-scale flow pattern over the North Pacific can be objectively characterized using the North Pacific Jet (NPJ) phase diagram.
- The NPJ phase diagram is constructed from the first two EOFs of Sept.–May 250-hPa zonal wind anomalies from the CFSR over the North Pacific (Fig. 1).




Figure 1: (a) Sept.–May 250-hPa mean zonal wind contoured in black every 10 m s⁻¹ above 30 m s⁻¹ and the regression of the first EOF to 250-hPa zonal wind anomaly data shaded in the fill pattern. (b) As in (a) but for the second EOF.




Figure 2: (a) 250-hPa wind speed shaded in the fill pattern at 0000 UTC 8 Nov 2014. (b) 250-hPa zonal wind anomalies at 0000 UTC 8 Nov 2014 projected onto the NPJ phase diagram.




Figure 3: (a) Centroids for 173 eastern U.S. extreme cold events during Sept.–May between 1979–2014 indicated by the red dots. (b) Average projection of 250-hPa zonal wind anomalies onto the NPJ phase diagram 3–7 days prior to each eastern U.S. extreme cold event in (a) identified with a red 'X'. The numbers in the blue boxes identify the number of extreme cold events associated with each jet regime.

Large-Scale Flow Evolution: 20 March 2016 – 3 April 2016




Figure 4: 500-hPa geopotential height contoured in blue every 6 dam, relative vorticity shaded in the fill pattern every 4 × 10⁻⁵ s⁻¹, and ascent contoured in green every 4 dPa s⁻¹ at 0000 UTC on the date listed in the top left of each panel.




Figure 5: The evolution of the flow pattern shown in Fig. 4 in terms of the NPJ phase diagram. Each diamond corresponds to 0000 UTC on the date listed immediately adjacent to the symbol. The green and red diamonds indicate the start and end of the trajectory, respectively.

- The 15-day period was characterized by discontinuous retrogression that culminated in the development of an omega block over the North Pacific on 28 March and a cold air outbreak over the Northeast U.S. on 3 April (Fig. 4).
- Throughout the period 20 March – 3 April, the NPJ was characterized by a jet extension and equatorward shift regime, which favors the development of below-normal temperatures over the eastern U.S. (Figs. 3 and 5).

Colorado Front Range Snowstorm






Figure 6: All plots show a series of GFS deterministic forecasts that verify at 1200 UTC 23 March 2016. The forecasted variables shown are the position of the (a) 552 dam geopotential height contour at 500 hPa, (b) 2 × 10⁻⁴ K s⁻¹ contour of 800–700-hPa layer-averaged warm air advection, (c) 132 dam geopotential height contour at 850 hPa, and (d) -1 Pa s⁻¹ contour of vertical motion at 500 hPa.



- Forecasts with lead times greater than 120 h generally positioned the 500-hPa trough too far west (Fig. 6a).
- A very subtle westward shift in the position of the 850-hPa geopotential height minimum and strongest 800–700-hPa warm air advection continued until 24 h prior to the event (Fig. 6b,c).
- The subtle westward shift in both the geopotential height minimum and the band of warm air advection had implications for where the strongest ascent would occur along the CO Front Range (Fig. 6d).

Omega Block Formation



- The 240-h GEFS forecast of 500-hPa geopotential height suggested a broad ridge over the eastern North Pacific and western North America, with considerable ensemble spread (Fig. 7a).
- The 120-h GEFS forecast showed strong agreement for the omega block over the North Pacific, but uncertainty remained with respect to a subsynoptic-scale trough over the MS River Valley (Fig. 7b).