

GEOLOGY OF THE BADGER BAY - SEAL BAY AREA, NORTH-CENTRAL NEWFOUNDLAND

by

K. Douglas Nelson

A Dissertation

Submitted to the State University of New York at Albany

in Partial Fulfillment of  
the Requirements for the Degree of  
Doctor of Philosophy

College of Science and Mathematics

Department of Geological Sciences

1979

State University of New York at Albany

COLLEGE OF ARTS AND SCIENCES

The dissertation submitted by

K. Douglas Nelson

under the title

GEOLOGY OF THE BADGER BAY - SEAL BAY AREA,

NORTH-CENTRAL NEWFOUNDLAND

has been read by the undersigned. It is hereby recommended for acceptance to the Faculty of the University in partial fulfillment of the requirements for the degree of Doctor of Philosophy.

(Signed)

(Signed)

(Signed)

(Signed)

(Date)

(Date)

(Date)

(Date)

Recommended by the Department of Geological Sciences

(Signed)

Kevin Burke, Chairman.

Recommendation accepted by the Dean of Graduate Studies for the Graduate Academic Council.

(Signed)

(Date)

GEOLOGY OF THE BADGER BAY - SEAL BAY AREA, NORTH-CENTRAL NEWFOUNDLAND

by

K. Douglas Nelson

Abstract of a Dissertation

Submitted to the State University of New York at Albany


in Partial Fulfillment of  
the Requirements for the Degree of  
Doctor of Philosophy

College of Science and Mathematics  
Department of Geological Sciences

1979

## ABSTRACT

Coastline in the Badger Bay - Seal Bay area of north-central Newfoundland exposes the thickest and least disrupted section of Ordovician rocks in Newfoundland's Central Volcanic Belt. The following conformable stratigraphic sequence is observed: 1) >5 km. of variegated mafic and silicic submarine volcanics and volcanoclastics of lower Ordovician age; 2) a thin (<.5 km.) sequence consisting of thin bedded red and green argillites, manganiferous cherts, bioturbated cherts and black sulferous graptolite-bearing argillites of Caradocian age; 3) >1.2 km. of quartz-rich sandstones of upper Ordovician age. Correlative sequences occur to the east in the Fortune Harbour Peninsula area and on New World Island. Together they record Early Ordovician island arc volcanism, Medial Ordovician cessation of volcanism and subsidence, and Medial through Late Ordovician uplift and erosion of a terrane to the north or west. Analysis of the detrital mineralogy and provenance of the Late Ordovician sediments indicates that they were derived from strata now exposed in the Burlington Peninsula area of western Newfoundland. This places an independent constraint on the age of Fleur de Lys metamorphism and deformation recorded in that area. It is concluded that ophiolite obduction on the Western Platform, Fleur de Lys metamorphism and deformation in the Burlington Peninsula area, and cessation of arc-type volcanism in the Notre Dame Bay area occurred synchronously. These results support the hypothesis that ophiolite obduction in western Newfoundland resulted from the collision of an island arc with an Atlantic-type continental margin during Medial Ordovician time.



"The Outcrops O.K."

# TABLE OF CONTENTS

	<u>Page</u>
ABSTRACT . . . . .	i
LIST OF FIGURES . . . . .	ii
LIST OF TABLES . . . . .	iv
LIST OF PLATES . . . . .	iv
ACKNOWLEDGMENTS . . . . .	v
INTRODUCTION . . . . .	1
General Statement . . . . .	1
Method of Approach . . . . .	3
<u>PART 1</u> <u>GEOLOGY OF THE BADGER BAY - SEAL BAY AREA</u>	
CHAPTER I	
FIELD METHODS . . . . .	7
CHAPTER II	
STRATIGRAPHY OF THE BADGER BAY - SEAL BAY AREA . . . . .	9
Introduction . . . . .	9
Wild Bight Group - Introduction . . . . .	12
Omega Point Formation . . . . .	14
Corner Point Formation . . . . .	22
Little Harbour Formation . . . . .	36
Seal Bay Head Formation . . . . .	46
Shoal Arm Formation . . . . .	54
Gull Island Formation . . . . .	60
CHAPTER III	
INTRUSIVE ROCKS . . . . .	70
Introduction . . . . .	70
Mafic Dikes, Sills and Plugs . . . . .	70
Lamprophyre Dikes . . . . .	73
Granitic Xenolith-Bearing Lamprophyre . . . . .	77
CHAPTER IV	
METAMORPHISM . . . . .	80
CHAPTER V	
STRUCTURE . . . . .	86
Regional Folds and Slaty Cleavage . . . . .	86
Faults . . . . .	90
Soft Sediment Folds . . . . .	92

## CHAPTER VI

BURTONS HEAD TERRANE . . . . .	96
Introduction . . . . .	96
Field Relations Around Burtons Head . . . . .	97
Field Relations Along the Southeast Shore of Sops Arm . . . . .	102
Sops Head Complex . . . . .	105
Intrusives in the Burtons Head Terrane . . . . .	120

PART 2  
REGIONAL GEOLOGY

## CHAPTER VII

SUMMARY OF EXPLOITS TERRANE STRATIGRAPHY IN NOTRE DAME BAY. . . . .	123
---	-----

## CHAPTER VIII

DETRITAL MINERALOGY AND PROVENANCE OF SANSOM-EQUIVALENT GREYWACKES . . . . .	132
Introduction . . . . .	132
Provenance . . . . .	132
Petrographic Description - Introduction . . . . .	134
Petrographic Description - Framework Grains . . . . .	135
Detrital Modes . . . . .	139
Conclusions . . . . .	148

## CHAPTER IX

CONCLUSIONS . . . . .	150
Regional Implications . . . . .	150
Problems - The Acadian Orogeny . . . . .	152
Problems - Strike-Slip Motions . . . . .	155

BIBLIOGRAPHY . . . . .	156
------------------------	-----

APPENDIX 1: Ophiolitic Detritus in the Upper Ordovician Flysch of Notre Dame Bay and its Bearing on the Tectonic Evolution of Western Newfoundland . . . . .	162
APPENDIX 2: The Age of the Roberts Arm Group, North-Central New- foundland: Discussion . . . . .	173
APPENDIX 3: Lower Paleozoic Trench-Fill Sequence New World Island, Newfoundland: Discussion . . . . .	178
APPENDIX 4: Fossil Localities in the Badger Bay - Seal Bay Area . . . . .	182

# LIST OF FIGURES

Figure Number		Page
1-1	Location of Badger Bay - Seal Bay area with respect to major tectonic zones in Newfoundland.	2
1-2	Major litho-stratigraphic subdivisions of the Central Volcanic Belt in the Notre Dame Bay area.	4
2-1	Stratigraphic nomenclatures for the Badger Bay - Seal Bay area, no relative thicknesses implied.	11
2-2	Stratigraphy of the Wild Bight Group and correlative pre-Caradocian strata in the adjacent New Bay - Fortune Harbour Peninsula area.	13
2-3	Location map, Omega Point Formation type section.	15
2-4	Fine-grained volcanoclastic sandstones and argillites, Omega Point Formation, shore south of Omega Point.	17
2-5	Slump folded horizon in Omega Point Formation, note undeformed beds above and below, shore south of Omega Point.	19
2-6	Isolated "amoeboid" shaped diabase body in red argillite, Omega Point Formation, shore south of Omega Point.	21
2-7	Location map, Corner Point Formation type section.	23
2-8	Purple silicic volcanic breccia, Corner Point Formation, west shore Seal Bay.	30
2-9	Sedimentary lamination(?) in massive grey felsite, Corner Point Formation, west shore Seal Bay.	30
2-10	Pillow lavas on westernmost island in bottom of Seal Bay, Corner Point Formation.	33
2-11	Location map, Little Harbour Formation type section.	37
2-12	Interbedded volcanoclastic sandstones and argillites, lower Little Harbour Formation, NW shore Side Harbour.	41
2-13	Volcanoclastic conglomerate, Little Harbour Formation, Gull Island in Seal Bay.	41
2-14	Location map, Seal Bay Head Formation type area.	47
2-15	Mafic volcanic breccia, Seal Bay Head Formation, Seal Bay Head.	51



	Page
2-16 Location map, Shoal Arm Formation type section.	55
2-17 Interbedded red and green volcanoclastic sandstones and argillites with thin manganiferous chert bed (purple), lower Shoal Arm Formation, SW end Gull Island in Badger Bay.	57
2-18 Thin bedded red argillites with steely blue manganese weathering stain, lower Shoal Arm Formation, SW end Gull Island.	57
2-19 Thin bedded grey bioturbated chert, middle Shoal Arm Formation, south shore Gull Island.	59
2-20 Graptolite bearing black slate, upper Shoal Arm Formation, south shore Gull Island.	59
2-21 Location map, Gull Island Formation type section.	61
2-22 71 reoriented flute casts, lower Gull Island Formation, NW shore Shoal Arm; outer circle = 20, arrows indicate direction of flute tails.	68
2-23 Interbedded greywacke and slate, lower Gull Island Formation, NW shore Shoal Arm.	65
2-24 Thick bedded greywacke, Gull Island Formation, NE end Gull Island.	65
3-1 Orientation of lamprophyre dikes near Leading Ticks.	74
3-2 Zenolith bearing lamprophyre dike, east shore Seal Bay just south of Wild Bight.	78
5-1 Northeast plunging mesoscopic fold in thin bedded volcanoclastic argillites, northwest shore Beaver Bight. Note well developed axial surface slaty cleavage.	87
5-2 "Nappe like" slump folds in volcanoclastic sandstones, east shore Seal Bay.	94
6-1 Schematic of mesoscopic structure in matrix of Sops Head Complex, Woody Island.	111
6-2 Regional geometry of mesoscopic fabric elements in the Sops Head - Boones Point Complex.	114
6-3 Large limestone block (olistolith?) surrounded by clastic sedimentary matrix, NW side Duck Island.	101
6-4 Mudstone conglomerate, Sops Head Complex, NW shore Sops Arm.	109

6-5	Limestone (grey) and "smeared" mafic volcanic (light grey green) in shaly matrix, Sops Head Complex on Green Island; trace of $S_1$ parallel to lower edge of photo, trace of $S_2$ parallel to pencil, pencil points toward small $F_2$ fold in smeared volcanic material.	116
6-6	$F_1$ isoclinal intrafolial fold hook in dismembered greywacke and slate, Sops Head Complex on north shore of Woody Island.	117
6-7	"Smeared" mafic volcanic material in shaly matrix, Sops Head Complex on Green Island.	118
7-1	Stratigraphy of the Exploits Terrane in Notre Dame Bay (Nelson, this study; Helwig, 1967; Kay, 1976).	124
7-2	Stratigraphic sequences on New World Island, after Kay (1976).	126
8-1	Summary map of paleocurrent data for Sansom-equivalent strata in the Notre Dame Bay area (1 this study; 2-7 Helwig, 1967; 8 (solid) Eastler, 1969).	133
8-2	A, Q-F-L plot of medium to coarse-grained sandstones from the New Bay Formation (pre-Caradocian); B, fields defined by both (neglecting NB-40).	142
8-3	Q-F-L comparison of Sansom-equivalent and New Bay Formation sandstones with sandstones of the Great Valley Sequence, California.	144

## LIST OF TABLES

## Table Number

8-1	Summary of point count results on Sansom-equivalent and New Bay Formation sandstones.	141
-----	---	-----

## LIST OF PLATES

Plate 1:	Coastline Geology of the Badger Bay - Seal Bay area.
Plate 2:	Generalized Geologic Map of the Notre Dame Bay area.
Plate 3:	Airphoto Lineament Map of the Badger Bay - Seal Bay area.

## ACKNOWLEDGMENTS

I gratefully acknowledge the advice, financial support, and editorial efforts provided by W.S.F. Kidd. Discussions with J. Casey, E. Rosencrantz, D. Rowley, J. Karson, J. Dewey, P. Dean, and H. Williams also contributed more or less to whatever understanding of Newfoundland geology I now have. The energy, patience, and relaxed good humor of John McCallen is gratefully acknowledged; this despite boredom, rain, bugs, celibacy, and innumerable "dusty" crossings. All of the residents of Cards Harbour, Newfoundland, deserve thanks for their unstinting hospitality, helpfulness, and good cheer. In particular, Ralph Lush gave me his boat for an entire summer, his father Gordon made sure I didn't drown, and Dott Henstridge filled me with so much fisherman's brewis that I surely would have if I had fallen in. Thanks to Caroline, Suzanne, Kathy, Marta M., Marta C., Frieda, Barb, Patti, Linda S., Linda U., Anne, Diane, and Donna - each contributed in her own special way. Special thanks goes to Marta Cary - for perspective. Linda Schroll obligingly and expertly typed the final draft of this dissertation.

This study was supported during the first year by Geological Society of America Penrose Grant No. 2234-77 awarded to the author, and during the latter two years by National Science Foundation Grant No. 320-832A awarded to W.S.F. Kidd.