

---

# An Examination of Low-Skill Arctic Cyclones During Summer

---

**Kevin Biernat, Lance Bosart, and Dan Keyser**  
*Department of Atmospheric and Environmental Sciences*  
*University at Albany, State University of New York*

Friday 26 February 2021  
ONR Arctic Cyclone DRI Meeting

Research Supported by ONR Grant N00014-18-1-2200

# Overview

---

- Identify periods of low and high forecast skill of the synoptic-scale flow over the Arctic and low-skill Arctic cyclones (ACs) occurring during these periods.
- Examine dynamical and thermodynamic quantities characterizing the Arctic environment and low-skill ACs during low-skill and high-skill periods.
- Conduct AC-centered composite analyses of intense low-skill ACs during low-skill periods to identify features and processes governing the evolution of these ACs.

# Arctic forecast skill evaluation

---






- Utilize day-5 forecasts of 500-hPa geopotential height initialized at 0000 UTC during summers (June, July, and August) of 2007–2017 from 11-member 1° GEFS reforecast dataset v2 (Hamill et al. 2013).
- Calculate area-averaged root mean square error (RMSE) of 500-hPa geopotential height over the Arctic, using ERA-Interim as verification.
- Calculated standardized anomaly of area-averaged RMSE ( $\sigma_{\text{RMSE}}$ ).

# Arctic forecast skill evaluation

---

- Refer to forecast days valid at day 5 associated with the top and bottom 10% of  $\sigma_{\text{RMSE}}$  as **low-skill days** and **high-skill days**, respectively.
- Refer to forecasts initialized 5 days prior to low-skill days and high-skill days as **low-skill forecasts** and **high-skill forecasts**, respectively.
- Refer to time periods through day 5 encompassed by low-skill forecasts and high-skill forecasts as **low-skill periods** and **high-skill periods**, respectively.

# Quantities characterizing the Arctic environment



**Legend**

Distribution of quantities area-averaged over the Arctic ( $\geq 70^\circ\text{N}$ )

- Low-skill mean
- High-skill mean
- 2007–2017 climo mean
- Statistically significant difference with respect to climo at 95% confidence level
- Statistically significant difference with respect to climo at 95% confidence level

**Shading:** Interquartile range      **Data:**  $1^\circ$  ERA-Interim

# Identification of low-skill ACs

---


- Create a 2007–2017 summer (June, July, and August) AC climatology by obtaining cyclone tracks from 1° ERA-Interim cyclone climatology prepared by Sprenger et al. (2017).
- Deem cyclones that last  $\geq 48$  h and spend at least some portion of their lifetimes in the Arctic ( $> 70^\circ\text{N}$ ) as ACs.

# Identification of low-skill ACs


---

- Track ACs in forecasts from GEFS reforecast dataset v2 by utilizing an objective sea level pressure (SLP)-based tracking algorithm (Crawford et al. 2020).
- Consider forecasts initialized 120 h prior to the time of lowest SLP of the ACs when located in the Arctic during low-skill and high-skill periods.
- Calculate 120-h intensity RMSE based on minimum SLP of the ACs at the aforementioned time of lowest SLP, using ERA-Interim as verification.
- Refer to ACs associated with the top 25% of 120-h intensity RMSE for low-skill and high-skill periods as low-skill ACs for these respective periods.

# Identification of low-skill ACs



## Legend



**Red:** Low-skill ACs during low-skill periods (N = 58)


**Blue:** Low-skill ACs during high-skill periods (N = 39)

○ mean

★ min and max



# Quantities characterizing low-skill ACs




## Legend

Distribution of most extreme value of quantities characterizing ACs when the ACs are located in the Arctic for the following categories:

**Gray:** All ACs in 2007–2017 climatology ( $N = 730$ )

**Red:** Low-skill ACs during low-skill periods ( $N = 58$ )

**Blue:** Low-skill ACs during high-skill periods ( $N = 39$ )



★ Statistically significant difference with respect to climatology at 95% confidence level

Area-averaged quantities are calculated within 1000 km of AC center

**Data:** 1° ERA-Interim

# AC-centered composites

---

- Composite top 25% strongest low-skill ACs during low-skill periods ( $N = 14$ ) at various lag times relative to the time of lowest SLP of the ACs when located in the Arctic using ERA5 ( $0.25^\circ \times 0.25^\circ$ ).


# AC-centered composites

---

- Composite top 25% strongest low-skill ACs during low-skill periods ( $N = 14$ ) at various lag times relative to the time of lowest SLP of the ACs when located in the Arctic using ERA5 ( $0.25^\circ \times 0.25^\circ$ ).
- For each lag time:
  - Determine mean latitude and longitude of ACs.

# AC-centered composites

- Composite top 25% strongest low-skill ACs during low-skill periods ( $N = 14$ ) at various lag times relative to the time of lowest SLP of the ACs when located in the Arctic using ERA5 ( $0.25^\circ \times 0.25^\circ$ ).
- For each lag time:
  - Determine mean latitude and longitude of ACs.
  - Rotate and project ERA5 grids to a  $25 \times 25$  km EASE2 equal area grid such that the AC center lies on y-axis ( $0^\circ$  longitude) of the EASE2 grid.




**EASE2 Grid**

(source: NSIDC: <https://nsidc.org/ease/ease-grid-projection-gt>)

# AC-centered composites


- Composite top 25% strongest low-skill ACs during low-skill periods ( $N = 14$ ) at various lag times relative to the time of lowest SLP of the ACs when located in the Arctic using ERA5 ( $0.25^\circ \times 0.25^\circ$ ).
- For each lag time:
  - Determine mean latitude and longitude of ACs.
  - Rotate and project ERA5 grids to a  $25 \times 25$  km EASE2 equal area grid such that the AC center lies on y-axis ( $0^\circ$  longitude) of the EASE2 grid.



SLP (black) on 25x25 km EASE2 grid (grid points in red) valid 1200 UTC 6 Aug 2012

# AC-centered composites


- Composite top 25% strongest low-skill ACs during low-skill periods ( $N = 14$ ) at various lag times relative to the time of lowest SLP of the ACs when located in the Arctic using ERA5 ( $0.25^\circ \times 0.25^\circ$ ).
- For each lag time:
  - Determine mean latitude and longitude of ACs.
  - Rotate and project ERA5 grids to a  $25 \times 25$  km EASE2 equal area grid such that the AC center lies on y-axis ( $0^\circ$  longitude) of the EASE2 grid.
  - Shift projected grids to mean latitude of ACs.




SLP (black) on 25x25 km EASE2 grid (grid points in red) valid 1200 UTC 6 Aug 2012

# AC-centered composites

Grid before shifting to mean latitude



Grid after shifting to mean latitude




# AC-centered composites

- Composite top 25% strongest low-skill ACs during low-skill periods ( $N = 14$ ) at various lag times relative to the time of lowest SLP of the ACs when located in the Arctic using ERA5 ( $0.25^\circ \times 0.25^\circ$ ).
- For each lag time:
  - Determine mean latitude and longitude of ACs.
  - Rotate and project ERA5 grids to a  $25 \times 25$  km EASE2 equal area grid such that the AC center lies on y-axis ( $0^\circ$  longitude) of the EASE2 grid.
  - Shift projected grids to mean latitude of ACs.
  - Rotate shifted grids to mean longitude of ACs.




# AC-centered composites




● AC location at lag 0 h (time of lowest SLP of AC in Arctic)


**Red lines** show tracks of ACs during lag -48 h to lag 36 h, when valid.



Time series of minimum SLP (hPa) of ACs (red) and of mean minimum SLP (hPa) of ACs (black) during lag -48 h to lag 36 h, when valid.




300-hPa wind speed ( $\text{m s}^{-1}$ )



● Mean AC location

— SLP (hPa)

--- 1000–500-hPa  
thickness (dam)




200-km area-averaged 500-hPa relative vorticity ( $10^{-5} \text{ s}^{-1}$ )




● Mean AC location

— 500-hPa geopotential height (dam)

▬ 500-hPa wind ( $\text{m s}^{-1}$ )




Integrated vapor transport (IVT) ( $\text{kg m}^{-1} \text{s}^{-1}$ )




● Mean AC location

→ IVT ( $\text{kg m}^{-1} \text{s}^{-1}$ )

— 700-hPa geopotential height (dam)



200-km area-averaged 350–250-hPa divergence ( $10^{-6} \text{ s}^{-1}$ )




● Mean AC location

→ 350–250-hPa  
irrotational wind ( $\text{m s}^{-1}$ )

— 350–250-hPa PV (PVU)

— Negative values of 800–600-hPa  $\omega$   
(every  $1 \times 10^{-3} \text{ hPa s}^{-1}$ )



850–600-hPa Eady growth rate ( $\text{day}^{-1}$ )



● Mean AC location

— SLP (hPa)

# Summary

---

- The Arctic environment tends to be characterized by more amplified synoptic-scale flow, greater baroclinic growth, and potentially greater latent heating during low-skill periods compared to high-skill periods.
- Low-skill ACs tend to be stronger and embedded in a region of more amplified synoptic-scale flow, greater baroclinic growth, and potentially greater latent heating during low-skill periods compared to high-skill periods.
- Intense low-skill ACs during low-skill periods intensify downstream of a mid-to-upper-tropospheric vortex in a region of relatively strong lower-to-midtropospheric baroclinicity, lower-to-midtropospheric ascent, tropospheric-integrated vapor transport, and upper-tropospheric divergence.
- A combination of baroclinic processes and latent heating likely play important roles in the intensification of intense-low-skill ACs during low-skill periods.