


Investigating the factors responsible for secondary eyewall formation in an ensemble of high-resolution hurricane simulations

Kristen L. Corbosiero and Ryan D. Torn
University at Albany


Motivation:

Secondary eyewalls are common in strong hurricanes




Motivation:



Eyewall replacement cycles are well understood



The formation of a secondary eyewall (SE) is followed by a known series of structural and intensity changes called an **eyewall replacement cycle (ERC)**.



Motivation: SE formation (SEF): Environmental moisture



Weak Gradient in Katrina

Strong Gradient in Rita

Rainbands

Eyewall




Dry

Moist

Ortt and Chen (2007)
Judt and Chen (2010)

Motivation:


SE formation (SEF): Environmental moisture



Weak Gradient in Katrina

Strong Gradient in Rita

Rainbands spiraling into the center from 100s of km away




Circular rainbands surrounding the center

Ortt and Chen (2007)
Judt and Chen (2010)



Model design: Advanced Hurricane WRF (AHW)

- ~ Run in **real time** at NCAR since 2004
- ~ **Three domains** with two-way moveable nests that follow the hurricane center
- ~ **AHW EnKF data assimilation** system generates a 96 member analysis ensemble every 6 h for a basin scale 36 km domain, and a 12 km TC-following nested domain
- ~ Analyses from **48 members** were used to initialize forecasts of **Igor** from 00 UTC 11 September out to 120 h




Results:

Track and intensity




Results:

SEF and ERC: Member #11



Results:


SEF and ERC: Member #11



Outward
propagating
features from
the eyewall


Results:

SEF and ERC: Member #11



Results:

SEF and ERC: Member #11




Moat
genesis
and SEF

Increase in
convective
activity

Outward
propagating
features from
the eyewall

Results:

SEF and ERC: Member #11




Inward movement and amplification

Increase in convective activity

Outward propagating features from the eyewall


Results:

SEF and ERC: Member #11



Results:

SEF and ERC: Member #11



Cessation of inner eyewall



Inward movement and amplification

Increase in convective activity

Outward propagating features from the eyewall


Results:

Comparison of two strong members




Results:

Comparison of two strong members





#35: Quick to develop; classic SEF & ERC between 60 h and 78 h



#30: Slow, outward movement of the eyewall and expansion of the convective field

Results:



Comparison of two strong members



#30 is drier and exhibits a stronger moisture gradient between the core and environment

Results:


Comparison of two strong members




#30 is drier and exhibits a stronger moisture gradient between the core and environment

Results: Composite evolution


10 m wind speed



200-850 hPa shear



0-850 hPa **upshear** PW




The members exhibiting **SEs** are more intense, especially in the time period after the ERCs are complete (78-102 h).


The time period of **SEF** featured below ensemble mean wind shear and significantly higher upshear PW.

Results: Composite evolution


10 m wind speed



200-850 hPa shear



0-850 hPa **upshear** PW




The members exhibiting **SEs** are more intense, especially in the time period after the ERCs are complete (78-102 h).


The time period of **SEF** featured below ensemble mean wind shear and significantly higher upshear PW.

Results: Composite evolution


10 m wind speed



200-850 hPa shear



0-850 hPa **upshear** PW




The members exhibiting **SEs** are more intense, especially in the time period after the ERCs are complete (78-102 h).


The time period of **SEF** featured below ensemble mean wind shear and significantly higher upshear PW.

Results: Composite evolution


10 m wind speed



200-850 hPa shear



0-850 hPa **upshear** PW




The members exhibiting **SEs** are more intense, especially in the time period after the ERCs are complete (78-102 h).

The time period of **SEF** featured below ensemble mean wind shear and significantly higher upshear PW.

Results:


Qvapor distribution: SEF members vs. ensemble mean



Results:

Qvapor distribution: SEF members vs. ensemble mean

Inner core
moistens, far
field **dries** as
storm spins up




Results:

Qvapor distribution: SEF members vs. ensemble mean

Inner core
moistens, far
field **dries** as
storm spins up

SEF members
are more
moist, but not
uniformly




Results:

Qvapor distribution: SEF members vs. ensemble mean

Inner core
moistens, far
field **dries** as
storm spins up


SEF members
are more
moist, but not
uniformly



Results:

Qvapor distribution: SEF members vs. ensemble mean


Inner core
moistens, far
field **dries** as
storm spins up



SEF members
are more
moist, but not
uniformly

Results:


Qvapor distribution: SEF members vs. ensemble mean



The largest Q_{vapor} anomalies at mid-levels are upshear and right of shear, while the outer rainband region is uniformly moist.

Conclusions:

Summary



- ~ The AHW EnKF system is able to realistically simulate the secondary eyewall formation and eyewall replacement cycle of Hurricane Igor (2010)
- ~ The SEF members are more intense, are embedded in environments with weaker shear, and are more moist
- ~ The outer rainband region is uniformly, anomalously moist, while enhanced moisture is predominantly upshear and right of shear in the core